



## POSITION STATEMENT OF THE NATIONAL LYMPHEDEMA NETWORK

*By NLN Medical Advisory Committee; Next Revision Date Spring 2011*

### TOPIC: LYMPHEDEMA RISK REDUCTION PRACTICES

- I. Skin Care - Avoid trauma / injury to reduce infection risk<sup>1,2</sup>**
  - Keep extremity clean and dry
  - Apply moisturizer daily to prevent chapping/chafing of skin<sup>2</sup>
  - Attention to nail care; do not cut cuticles
  - Protect exposed skin with sunscreen and insect repellent
  - Use care with razors to avoid nicks and skin irritation
  - If possible, avoid punctures such as injections and blood draws
  - Wear gloves while doing activities that may cause skin injury (i.e., washing dishes, gardening, working with tools, using chemicals such as detergent)
  - If scratches/punctures to skin occur, wash with soap and water, apply antibiotics, and observe for signs of infection (i.e. redness)
  - If a rash, itching, redness, pain, increased skin temperature, fever or flu-like symptoms occur, contact your physician immediately for early treatment of possible infection<sup>3</sup>
  
- II. Activity / Lifestyle**
  - Gradually build up the duration and intensity of any activity or exercise<sup>4-8</sup>
  - Take frequent rest periods during activity to allow for limb recovery
  - Monitor the extremity during and after activity for any change in size, shape, tissue, texture, soreness, heaviness or firmness<sup>9</sup>
  - Maintain optimal weight<sup>10-13</sup>
  
- III. Avoid Limb Constriction<sup>9</sup>**
  - If possible, avoid having blood pressure taken on the at-risk extremity
  - Wear loose fitting jewelry and clothing
  
- IV. Compression Garments<sup>2,9,14</sup>**
  - Should be well-fitting
  - Support the at-risk limb with a compression garment for strenuous activity (i.e. weight lifting, prolonged standing, running) except in patients with open wounds or with poor circulation in the at-risk limb<sup>15</sup>
  - Consider wearing a well-fitting compression garment for air travel<sup>16,17</sup>

## **V. Extremes of Temperature<sup>1</sup>**

- Avoid exposure to extreme cold, which can be associated with rebound swelling, or chapping of skin
- Avoid prolonged (greater than 15 minutes) exposure to heat, particularly hot tubs and saunas
- Avoid placing limb in water temperatures above 102°Fahrenheit (38.9°Celsius)

## **VI. Additional Practices Specific to Lower Extremity Lymphedema**

- Avoid prolonged standing, sitting<sup>18</sup> or crossing legs
- Wear proper, well-fitting footwear and hosiery<sup>19,20</sup>
- Support the at-risk limb with a compression garment for strenuous activity except in patients with open wounds or with poor circulation in the at-risk limb<sup>15</sup>

NOTE: Given that there is little evidence-based literature regarding many of these practices, the majority of the recommendations must at this time be based on the knowledge of pathophysiology and decades of clinical experience by experts in the field.

## References

1. Rich A: How to care for uncomplicated skin and keep it free of complications. *Br J Community Nurs* 12:S6-9, 2007
2. Zuther JE: *Lymphedema Management: The Comprehensive Guide for Practitioners*. New York, NY, Thieme Medical Publishers, Inc., 2005
3. Feldman JL: The challenge of infection in lymphedema. National Lymphedema Network LYMPH Link Article Reprint, 2005
4. Johansson K, Tibe K, Weibull A, et al: Low intensity resistance exercise for breast cancer patients with arm lymphedema with or without compression sleeve. *Lymphology* 38:167-80, 2005
5. Moseley AL, Piller NB, Carati CJ: The effect of gentle arm exercise and deep breathing on secondary arm lymphedema. *Cancer Control* 38:136-145, 2005
6. Ahmed RL, Thomas W, Yee D, et al: Randomized controlled trial of weight training and lymphedema in breast cancer survivors. *J Clin Oncol* 24:2765-72, 2006
7. Bicego D, Brown K, Ruddick M, et al: Exercise for women with or at risk for breast cancer-related lymphedema. *Phys Ther* 86:1398-405, 2006
8. Lasinski BB: Exercise, lymphedema and the limb at risk. National Lymphedema Network LYMPH Link Article Reprint, 2001
9. Moffatt CJ: Best practice for the management of lymphoedema. *Lymphoedema Framework International Consensus*. London, MEP Ltd, 2006
10. Werner RS, McCormick B, Petrek J, et al: Arm edema in conservatively managed breast cancer: Obesity is a major predictive factor. *Radiology* 180:177-84, 1991
11. Goffman TE, Laronga C, Wilson L, et al: Lymphedema of the arm and breast in irradiated breast cancer patients: risks in an era of dramatically changing axillary surgery. *Breast J* 10:405-11, 2004
12. Shaw C, Mortimer P, Judd PA: Randomized controlled trial comparing a low-fat diet with a weight-reduction diet in breast cancer-related lymphedema. *Cancer* 109:1949-56, 2007

13. Ridner SH, Dietrich MS: Self-reported comorbid conditions and medication usage in breast cancer survivors with and without lymphedema. *Oncol Nurs Forum* 35:57-63, 2008
14. Stewart PJ: The garments we wear. National Lymphedema Network LymphLink, 2007
15. Marston W, Vowden K: Understanding compression therapy. In European Wound Management Association (EWMA). *Compression therapy: A guide to safe practice*. London: MEP Ltd, 2003, pp 11-17
16. Casley-Smith JR, Casley-Smith JR: Aircraft flights and SCUBA diving. National Lymphedema Network Newsletter Article Reprint, 1995
17. Casley-Smith JR: Tips for travel and aircraft flight, National Lymphedema Network LYMPH Link Article Reprint, 2000
18. Ryan TJ: Risk factors for the swollen ankle and their management at low cost: Not forgetting lymphedema. *Int J Low Extrem Wounds* 1:202-8, 2002
19. Hewitson J: Foot care for lower extremity lymphedema. National Lymphedema Network LYMPH Link Article Reprint, 2001
20. White J: Lymphedema and your feet, National Lymphedema Network Newsletter Article Reprint, 1991
21. Földi M, Földi E, Kubik S: Prevention, *Textbook of Lymphology for Physicians and Lymphedema Therapists* (ed 5th German). Urban & Fischer, 2003, pp 275-279

© 2008 National Lymphedema Network (NLN). Permission to duplicate this handout as-is, in its entirety, for educational purposes only, not for sale. All other rights reserved. For reprint permission, please contact the NLN office at 510-208-3200 or [nln@lymphnet.org](mailto:nln@lymphnet.org).